

Ana Filipa da Silva Ferreira

Uma revisão sistemática sobre monitorização terapêutica do
infliximab e adalimumab: níveis, resultados clínicos e ensaios

A systematic review on infliximab and adalimumab drug
monitoring: levels, clinical outcomes and assays

março, 2017

Ana Filipa da Silva Ferreira

Uma revisão sistemática sobre monitorização terapêutica do
infliximab e adalimumab: níveis, resultados clínicos e ensaios

A systematic review on infliximab and adalimumab drug
monitoring: levels, clinical outcomes and assays

Mestrado Integrado em Medicina

Área: Farmacologia

Tipologia: Monografia

Trabalho efetuado sob a Orientação de:

Professor Doutor Fernando Magro

Trabalho organizado de acordo com as normas da revista:

Inflammatory Bowel Diseases

março, 2017

FMUP

Eu, Ana Filipa da Silva Ferreira, abaixo assinado, nº mecanográfico 201104870, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 11 / 03 / 2017

Assinatura conforme cartão de identificação:

Ana Filipa da Silva Ferreira

Projecto de Opção do 6º ano – DECLARAÇÃO DE REPRODUÇÃO

NOME

Ana Filipa da Silva Ferreira

NÚMERO DE ESTUDANTE

201104870

DATA DE CONCLUSÃO

22/03/2017

DESIGNAÇÃO DA ÁREA DO PROJECTO

Farmacologia

TÍTULO DISSERTAÇÃO/MONOGRAFIA (riscar o que não interessa)

A Systematic Review on Infiximab and Adalimumab Drug Monitoring: Levels, Clinical Outcomes and Assays

ORIENTADOR

Professor Doutor Fernando Magro

COORIENTADOR (se aplicável)

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTES TRABALHOS APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTES TRABALHOS (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input checked="" type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTES TRABALHOS.	<input type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 11 / 03 / 2017

Assinatura conforme cartão de identificação:

Ana Filipa da Silva Ferreira

A Systematic Review on Infliximab and Adalimumab Drug Monitoring: Levels, Clinical Outcomes and Assays

Filipa Silva-Ferreira, MD,^{*,†} Joana Afonso, MSc,^{*,†} Pedro Pinto-Lopes, MD,[‡]
and Fernando Magro, MD, PhD^{*,†,§} on behalf of GEDII (Portuguese IBD Study Group)

Background: Immunogenicity to therapeutic proteins has been linked to loss of response by a large percentage of patients taking anti-tumor necrosis factor- α agents. Drug monitoring can be extremely useful, allowing physicians to adjust the therapeutic scheme individually. This article aims to systematically review the published data with respect to cutoff levels of infliximab (IFX) and adalimumab (ADA) and relate them to the methodology adopted for quantification of IFX and ADA levels and clinical outcomes.

Methods: The PubMed database was searched to identify studies focusing on the association between IFX or ADA cutoff levels and clinical outcomes in patients with inflammatory bowel disease.

Results: Of the 1654 articles initially selected by queries, 20 were included. A receiver operating characteristic curve analysis was performed to identify cutoff levels of IFX or ADA that correlated with a clinical outcome, but only 6 studies performed the same analysis for antidrug antibody levels. Cutoff levels were different between studies. The methodology chosen for level quantifications, clinical outcomes, and sample size and characteristics were also different. Nevertheless, measurement of drug levels should be performed during maintenance, and with loss of response, with persistent high levels of C-reactive protein, and when mucosal lesions are still present. In these scenarios, drug and antidrug levels were correlated with clinical outcomes.

Conclusions: Concerning drug levels monitoring any methodology is adequate. With respect to antidrug antibody levels, it will be necessary to define a gold standard method or to establish different cutoff levels for different methodologies.

(*Inflamm Bowel Dis* 2016;0:1–13)

Key Words: anti-infliximab antibodies, clinical outcomes, infliximab trough levels, therapeutic drug monitoring

Infliximab (IFX) and adalimumab (ADA) are antitumor necrosis factor- α (TNF α) agents that have changed the clinical course of many autoimmune diseases such as inflammatory bowel disease (IBD), psoriasis, and rheumatoid arthritis. These agents have been successfully used in the past decades to treat patients

with IBD, even in those who were refractory to conventional therapy.^{1–5} Introduction of these agents to the drug market allowed physicians to aim for more than clinical remission, as these new drugs were proven to induce endoscopic remission and mucosa healing in patients with either Crohn's disease (CD) or ulcerative colitis (UC).^{6–8} Despite this, up to 70% of patients lose responsiveness over time.⁹ Many mechanisms may be involved in the loss of response, but immunogenicity to the antibody itself is so far the best studied.¹⁰ The presence of antibodies to IFX (ATIs) in patients' serum was associated with a 3-fold higher risk of loss of response than in patients who did not have ATIs in their serum.⁹ Although ADA is a fully human monoclonal antibody drug, immunogenicity to this drug has already been described and a negative correlation between the presence of antibodies to ADA (ATA) and ADA trough levels (TLs) was demonstrated.¹¹ However, the influence of ADA levels in clinical and endoscopic remission is not well established yet.

When patients lose response to anti-TNF α agents, their physicians have roughly 4 options: (1) dose escalation, (2) addition of an immunomodulator, (3) change to another class of drugs, or (4) change to another anti-TNF agent.^{12–17} Currently, physicians have to empirically decide since measurement of drug and antidrug antibody levels is not yet used in daily practice. Many authors have highlighted the importance of knowing drug and antidrug antibody levels to better adjust the therapeutic scheme.

Received for publication February 26, 2016; Accepted April 25, 2016.

From the *Department of Pharmacology and Therapeutics, Faculty of Medicine, University of Porto, Porto, Portugal; †MedInUP, Center for Drug Discovery and Innovative Medicines, University of Porto, Porto, Portugal; and ‡Departments of Internal Medicine, and §Gastroenterology, Faculty of Medicine, Centro Hospitalar S. João, Porto, Portugal

Supported by the Portuguese Group of Studies in Inflammatory Bowel Disease (GEDII), Portugal, and partially by the European Regional Development Fund (ERDF); COMPETE (Operational Programme Thematic Factors of Competitiveness) and Fundação para a Ciência e a Tecnologia, Portugal (FCT) [grant number EXPL/IVC-PEC/1302/2013].

The authors have no conflict of interest to disclose.

F. S. Ferreira and J. Afonso contributed equally to the article. F. S. Ferreira and J. Afonso were involved in data collection and interpretation, and drafting the manuscript. P. Pinto-Lopes was involved in the data interpretation. F. Magro was involved in the conception and design of the study, interpretation of data, and drafting and revision of the manuscript. All authors read and approved the final manuscript.

Address correspondence to: Fernando Magro, MD, PhD, Department of Pharmacology and Therapeutics, Faculty of Medicine, University of Porto, Rua Plácido Costa, 4200 Porto, Portugal (e-mail: fm@med.up.pt).

Copyright © 2016 Crohn's & Colitis Foundation of America, Inc.

DOI 10.1097/MIB.0000000000000855

Published online.

Nonetheless, most authors emphasize the need to find a valid assay, especially to measure antidrug antibodies and to set cutoff levels to help in decision-making.^{18–20} The aim of this article was to systematically review the published data with respect to IFX and ADA levels, the methodology applied, and the relationship with clinical outcomes.

MATERIALS AND METHODS

A systematic review focusing on the association between IFX, ADA TL, ATIs, ATAs, and clinical outcomes in patients with IBD was performed.

Search Strategy

A literature search was performed, through July 2015, using the PubMed database with the following keywords and Medical Subject Headings (MeSH) terms: “{(adalimumab[All fields]) OR (infliximab[All fields])} AND {(inflammatory bowel disease [MeSH Terms]) OR (inflammatory bowel diseases[MeSH Terms]) OR (crohn’s disease[MeSH Terms]) OR (colitis, ulcerative[MeSH Terms]) OR (crohn disease[MeSH Terms])} AND ([clinical response] OR [clinical remission] OR [disease activity] OR [clinical outcomes]).” Considering this is a hot topic, we decided, on December 2015, to perform an additional literature search on abstracts presented on 3 reference congresses. The European Crohn’s and Colitis Organisation (ECCO) Website was searched for all published abstracts related with this topic, using the terms “infliximab ifx” and “adalimumab ada”; The United European Gastroenterology Week (UEGW) Website was searched for abstracts from the last United European Gastroenterology week in Barcelona; The Digestive Disease Week (DDW) Website was searched for abstracts from the past 5 years, using the terms “infliximab levels” and “adalimumab levels” in title, abstract, or keywords.

Eligibility Criteria

The inclusion criteria were: (1) articles studying the association between IFX or ADA cutoff levels and clinical outcomes in patients with IBD and (2) articles written in English.

We excluded studies that (1) were systematic reviews, (2) used another anti-TNF- α agent rather than IFX or ADA, (3) enrolled patients with other diseases rather than IBD (psoriasis, rheumatoid arthritis), (4) only assessed the relationship between IFX or ADA TL and clinical outcomes but did not perform a receiver operating characteristic (ROC) curve analysis, or (5) did not present the specificity and sensitivity values of the ROC curve analysis. This last criterion was defined so that we could infer the accuracy of the cutoff value (i.e., a cutoff value with a sensitivity and/or specificity of 50% would be no better at identifying true positives than flipping a coin). It was not applied to abstracts found on ECCO, UEGW, or DDW databases.

Study Selection and Data Collection Process

Studies were screened and selected by 2 reviewers. First, all titles and abstracts were read and the inclusion and exclusion

criteria were applied. Second, the articles considered for inclusion after selection by title/abstract reading were read fully and the inclusion and exclusion criteria were applied again. The data collected from each study were: the type of study and location, number of patients enrolled, and the type of IBD, definitions of clinical outcomes, antidrug antibodies incidence, type of assay used to measure IFX/ADA and ATIs/ATAs serum levels, and the results from the ROC curve analysis (cutoff levels and specificity and sensitivity values), except for the studies obtained in ECCO, UEGW, or DDW databases. In these studies, we have only had access to the abstract. A quality assessment was performed using a qualitative classification of the risk of bias. We used a 4-item classification based on the Meta-analysis of Observational Studies in Epidemiology checklist.²¹ The items were chosen based on the factors that can incorporate bias, i.e., inclusion and exclusion criteria, justification of the cohort (eligibility criteria, sources and methods of selecting participants, and the methods used to describe follow-up), the type of disease (if they pointed out whether the patients included had CD or UC), and the assay used to measure drug and antidrug antibody levels (Fig. 1).

RESULTS

Search and Study Selection

A total of 1237 articles were identified with our query (Fig. 2). Of these, 1160 were excluded by title and/or abstract alone, mainly because they did not study the association between IFX or ADA TL and clinical outcomes. Therefore, 77 articles were considered for full text analysis and after that 13 were included in our systematic review (Fig. 2). Two additional articles were included after searching those related to the 13 articles selected by query.^{22,23} From the search on ECCO, UEGW, and DDW abstract databases, 417 abstracts were found but only 5 were included, according to the inclusion criteria previously defined (Fig. 3).

Description of Studies

Of the 20 studies included, all but one²⁴ were conducted in adult patients. One study²⁵ only involved patients with UC, 11 studies^{6,11,22,23,26–33} only encompassed patients with CD and 7 studies pertained to patients with either UC or CD.^{8,24,34–38} Fifteen of the 19 studies involved IFX maintenance therapy^{6,8,22,24–28,30–35,37} (Table 1), whereas the other 4 involved ADA maintenance therapy (Table 2).^{11,23,29,36} One study encompassed patients from both regimens, IFX and ADA maintenance therapy.³⁸ Seven studies did not report information about the incidence of ATIs,^{25,29,31–33,38,39} and only 6 performed an ROC curve analysis to find a cutoff value for ATI^{22,32,34,35,37} or ATA¹¹ levels.

In 6 studies, the clinical outcome was “clinical remission,”^{8,11,22,24,28,36} usually assessed by the Harvey-Bradshaw Index–Mayo score and/or C-reactive protein (CRP) levels. In 4 studies, the outcome was “loss of response,”^{26,32,35,37} defined as an initial good clinical response to IFX induction treatment followed by a loss of clinical response to IFX during maintenance treatment

	Inclusion and exclusion criteria	Justification cohort	Type of IBD	Assay used
Echarri et al. 2015 ²⁸	?	?	+	+
Roblin et al. 2015 ³²	?	?	+	+
Ungar et al. 2015 ³⁸	?	?	?	?
Adedokun et al. 2014 ²⁵	+	+	+	+
Cornillie et al. 2014 ²⁷	+	+	+	+
Levesque et al. 2014 ³⁰	+	+	+	+
Marits et al. 2014 ⁸	-	-	+	+
Papamichail et al. 2015 ³¹	?	+	+	+
Singh et al. 2014 ²⁴	-	+	+	+
Tang et al. 2014 ³³	?	?	+	?
Vande Casteele et al. 2014 ²²	+	+	+	+
Bortlik et al. 2013 ²⁶	+	+	+	+
Imaeda et al. 2014 ⁶	+	+	+	+
Paul et al. 2013 ³⁷	+	+	+	+
Vande Casteele et al. 2013 ³⁴	+	+	+	+
Steenholdt et al. 2011 ³⁵	+	+	+	+
Zittan et al. 2016 ²³	+	+	+	+
Mazor et al. 2014 ¹¹	+	+	+	+
Roblin et al. 2014 ³⁶	-	+	+	+
Imaeda et al. 2013 ²⁹	-	+	+	+

FIGURE 1. Summary of risk of bias.




FIGURE 2. Data collection process.

leading to discontinuation of the drug. For Adedokun et al,²⁵ the endpoint was the “clinical response” defined as a decrease from the baseline in the total Mayo score of ≥ 3 points and at least 30%, and a decrease in the subscore for rectal bleeding of ≥ 1 or an absolute subscore for rectal bleeding of 0 or 1. For Levesque et al,³⁰ there were 2 endpoints which were an “increase in CD activity index ≥ 70 ” and an “increase in CRP ≥ 5 mg/L.” Imaeda et al⁶ defined 2 endpoints for IFX, including “mucosa healing,” meaning an endoscopic score of 0 or 1, and “CRP ≤ 0.3 mg/L,” whereas for ADA,²⁹ they only used “CRP ≤ 0.3 mg/L.” Four more studies defined “mucosa healing” as the endpoint of interest.^{23,31,33,38} Cornillie et al²⁷ defined clinical outcome as a “sustained response at week 54,” which was expressed as clinical remission based on the relevant disease activity index at week 54, in the absence of any dose intensification during IFX maintenance therapy. Paul et al³⁷ also defined 2 endpoints: “loss of response” and “absence of clinical remission.” Vande Casteele et al³⁴ described 3 endpoints which were “ATI formation,” “IFX discontinuation,” and “unsuccessful intervention.” The intervention (change in therapy) was considered successful if, at the second infusion after the intervention, the symptoms had disappeared and CRP, if elevated before the intervention, had




FIGURE 3. Data collection process.

TABLE 1. IFX Trough Levels and Antidrug Antibodies Cutoff, Methodology and Clinical Outcomes

Authors	Study Design	Population	Regimen	Country	Time Point
Echarri et al ²⁸	—	36 Adults with CD	IFX	Spain	W0, W6, W14, W30
Roblin et al ³²	Prospective cohort	119 Adults with CD	IFX	France	Trough level
Ungar et al ³⁸	Retrospective cross-sectional	78 Adults with IBD	IFX	Israel	No data
Adedokun et al, 2014 ²⁵	Observational (post-hoc ACT1-2)	454 Adults with UC	IFX; induction regimen followed by maintenance therapy	globally	W8 W30 W54 W14
Cornillie et al ²⁷	Observational (analyses of ACCENT I)	573 Adults with CD	IFX; induction regimen followed by maintenance therapy	North America, Europe, Israel	W8 Just before next infusion
Levesque et al ³⁰	Prospective cohort	327 Adults with CD	IFX; maintenance therapy	Canada	W8
Marits et al ⁸	Retrospective	63 Adults with CD, 15 adults with UC, 1 adult with U-IBD	IFX	Sweden	Just before next infusion
Papamichail et al, 2015 ³¹	Retrospective	101 adults with CD	IFX	Belgium	W 0, 2, 6, 14
Singh et al ²⁴	Prospective cohort	58 pediatric patients (<21 years) with CD and UC	IFX; induction regimen followed by maintenance therapy	USA	W14
Tang et al ³³	No	15 adults with CD	IFX	China	No data
Vande Casteele et al ²²	Observational	483 adults with CD	IFX; maintenance therapy	Belgium Canada	No data
Bortlik et al ²⁶	Retrospective	84 adults with CD	IFX	Czech Republic	W14–22
Imaeda et al, 2014 ⁶	Prospective cohort	65 adults with CD	IFX; maintenance therapy	Japan	Just before next infusion
Paul et al ³⁷	Prospective cohort	103 adults with IBD	IFX; maintenance therapy	France	Just before next infusion
Vande Casteele et al ³⁴	Retrospective	64 adults with CD, 26 adults with UC	IFX	Belgium	Just before next infusion
Steenholdt et al ³⁵	Retrospective	85 adults with CD, 21 adults with UC	IFX	Denmark	Just before next infusion

Authors	Drug			Antidrug antibodies				Endpoint
	Method	Cutoff, µg/mL	Spec/Sens, %	Method	Incidence, n (%)	Cutoff	Spec/Sens, %	
Echarri et al ²⁸	ELISA	>3 (w6)	No	ELISA	No (26)	No	No	Good response and sustained remission
Roblin et al ³²	ELISA (commercial kit, Theradiag)	<2	No	ELISA	No	>20 ng/mL	No	Loss of response
Ungar et al ³⁸	No	>5	85/—	No	No	No	No	Mucosa healing

TABLE 1 (Continued)

Authors	Drug			Antidrug antibodies				Endpoint
	Method	Cutoff, $\mu\text{g/mL}$	Spec/Sens, %	Method	Incidence, n (%)	Cutoff	Spec/Sens, %	
Adedokun et al, 2014 ²⁵	Classic ELISA	>41 (w8) >3.7 (w30) >1.7 (w54)	62/63 (w8) 71/65 (w30) 64/89 (w54)	Bridging ELISA	No	No	No	Clinical response
Cornillie et al ²⁷	Classic ELISA	≥ 3.5	78/64	Bridging ELISA	ATI+ = 2 (9)	No	No	Sustained response at w54
Levesque et al ³⁰	HMSA (commercial kit, Prometheus Laboratories)	≤ 2.8 –4.6 (a) ≤ 2.7 –2.8 (b)	68/61 (a) 74/64 (b)	HMSA (commercial kit)	ATI+ = 57 (18)	No	No	(a) Increased CDAI ≥ 70 ; (b) Increased CRP ≥ 5 mg/L
Marits et al ¹⁸	Classic ELISA	>4.1 (CD)	44/87	Inhibition ELISA	ATI+ = 22 (79); ATI transient = 4 (18)	No	No	Remission (HBI-Mayo and CRP)
Papamichail et al, 2015 ³¹	ELISA	>22.5 (w2) >12.8 (w6)	No	No	No	No	No	Short-term mucosa healing
Singh et al ²⁴	Classic ELISA and HMSA (Prometheus Laboratories)	≥ 5 ≥ 7	85/50 100/33	Bridging ELISA and HMSA (Prometheus Laboratories)	No (10 at w14), no (26 at w54)	No	No	Week 54 persistent remission
Tang et al ³³	No	>4.87	77/88	No	No	No	No	Mucosa healing
Vande Casteele et al ²²	HMSA (commercial kit)	>2.79	77.6/52.5	HMSA (commercial kit)	23.7% (IFX–/ATI– = 6.5%; IFX +/ATI– = 69.8%; IFX–/ATI+ = 16.4%; IFX+/ATI+ = 7.3%)	< 3.15 U/mL	87.4/38.0	Remission (CRP ≤ 5 mg/L)
Bortlik et al ²⁶	Classic ELISA (Q-INFLIXI, Matriks Biotek)	<3	62/70	Bridging ELISA (commercial kit)	ATI+ = 14 (17), ATI– = 24 (28), ATI inconclusive = 46 (55)	No	No	Loss of response at 1 year using IFX
Imaeda et al, 2014 ⁶	Classic ELISA	>4 (a) >0.6 (b)	70/71 (a) 62/73 (b)	ELISA (+acid dissociation and immunoaffinity chromatography)	No	No	No	(a) Mucosa healing (b) CRP < 0.3 mg/L
Paul et al ³⁷	ELISA (commercial kit, Theradiag)	<2 (a)	82.3/76	Bridging ELISA (commercial kit)	ATI+ = 34 (32.8), CD = 25.4%, UC = 41.5%, ^a ATI+ = 65.3% ^b	>200 ng/mL (b)	93.5/22.0	(a) Absence of clinical remission (b) Loss of response.

TABLE 1 (Continued)

Authors	Drug		Antidrug antibodies			
	Method	Cutoff, $\mu\text{g/mL}$	Spec/Sens, %	Method	Incidence, n (%)	Cutoff
Vande Casteele et al ³⁴	Classic ELISA	<13 (w6) (a)	81/72	HMSA	ATI+ = 53 (59), Transient ATI = 15 (28), Sustained ATI = 38 (72)	>9.1 U/mL (b)
		<2.2 (w14) (a)	94/79			
		<2.2 (w14) (c)	74/82			
Steenholdt et al ³⁵	Fluid-phase RIA	<0.5 (CD)	85/86 (CD)	Fluid-phase RIA	ATI+ = 35 (33.3)	≥ 10 U/mL (CD and UC)
		<0.8 (UC)	100/75 (UC)			

^aCutoff for ATI positivity > 10 ng/mL.^bCutoff for ATI positivity > 5 ng/mL.

CDAL, CD activity index; HBI, Harvey-Bradshaw index; sens, sensitivity; spec, specificity; U-IBD, unclassified inflammatory bowel disease.

decreased by >50% than the value at the time of loss of clinical response. Quality assessment was limited in those cases to which we only had access to the abstract.^{28,31–33,38} Taking into consideration the other studies, all but 4 had suitable inclusion and exclusion criteria,^{8,24,29,36} and all papers indicated the type of IBD and the assay used to measure drug and antidrug antibody levels.

Assays Used to Measure Drug and Antidrug Antibody Levels

One aspect that should be taken into consideration when analyzing drug TL and antidrug antibody levels is the assay used to measure them. All but 3 of the included works measured IFX or ADA TL using classic enzyme-linked immunosorbent assay (ELISA).^{6,8,11,24–29,31,32,34,36,37} Zittan et al,²³ Levesque et al,³⁰ and Vande Casteele et al²² used a homogeneous mobility shift assay (HMSA), whereas Steenholdt et al³⁵ used a fluid-phase radioimmunoassay (fluid-phase RIA). Singh et al²⁴ tested 2 methodologies, the classic ELISA and HMSA.

Regarding antidrug antibody measurements, 7 studies used bridging ELISA, either via home-made assays or commercial kits.^{25–28,32,36,37} Mazor et al¹¹ applied an adaptation of the anti-human lambda chain-based ELISA. In 2 studies by Imaeda et al, ATI⁶ and ATA²⁹ levels were also measured with ELISA, but samples were previously treated with acid in order to dissociate immune complexes. Other methods were used, namely HMSA,²³ fluid-phase RIA,³⁵ and inhibition ELISA.⁸ Singh et al²⁴ tested 2 methodologies, bridging ELISA and HMSA. Figure 4 displays all methodologies used.

Infliximab Levels

By Week of Measurement

Of the 16 IFX studies, 7 specified the time point measurement^{24–27,30}; one measured drug levels at week 2,³¹ 2 at week 6,^{28,31} 2 at week 8,^{25,30} 3 at week 14,^{24,26,27} 1 at week 22,²⁶ and 1²⁵ also measured IFX levels at weeks 30 and 54. Others only indicated that measurements were made before each infusion, thus representing drug TL.^{6,8,11,29,34,35,37}

In Papamichael et al,³¹ 2 cutoff levels were proposed (Table 1), both correlating with short-term mucosa healing, but after multiple logistic regression analysis, only IFX levels >12.8 $\mu\text{g/mL}$ at week 6 were retained as an independent factor to predict short-term mucosa healing (OR: 3.6, $P = 0.004$). Echarri et al²⁸ presented a largely different cutoff level for the same time point. They suggest that IFX levels >3 $\mu\text{g/mL}$ at week 6 had a positive-predictive value for “good response and sustained remission” of >90%. Adedokun et al²⁵ showed that IFX levels >41 $\mu\text{g/mL}$ at week 8 correlated with clinical response with a specificity of 62% and a sensitivity of 63% (Table 1). The median serum IFX concentration was significantly higher at week 8 in patients with clinical response or mucosal healing during induction than those not achieving these endpoints. Levesque et al³⁰ found a different cutoff: a mean IFX trough concentration <3 $\mu\text{g/mL}$ at week 8 was

TABLE 2. ADA Trough Levels and Antidrug Antibodies Cutoff, Methodology and Clinical Outcomes

Authors	Study Design	Population	Regimen	Country	Time point	Drug		
						Method	Cutoff, $\mu\text{g/mL}$	Spec/Sens, %
Zittan et al, 2016 ²³	Observational	60 Adults with CD	ADA	Canada	—	HMSA (commercial kit, Prometheus Laboratories)	8.14	76.0/91.4
Ungar et al ³⁸	Retrospective, cross-sectional	67 Adults with IBD	ADA	Israel	—	—	>7.1	85/—
Mazor et al ¹¹	Observational, cross-sectional	71 Adults with CD	ADA	Israel	Just before next infusion	Classic ELISA	>5.85 (a)	70.6/68
Roblin et al ³⁶	Observational cross-sectional	40 Adults with IBD	ADA; maintenance therapy	France	W22	ELISA (commercial kit, Theradiag)	<4.9 (a) >4.85 (b)	85/66 67/81
Imaeda et al, 2014 ²⁹	Prospective cohort	40 Adults with CD	ADA; maintenance therapy	Japan	Just before next infusion	Classic ELISA	>5.9	92/67

Authors	Antidrug antibodies				Endpoint
	Method	Incidence, n (%)	Cutoff	Spec/Sens, %	
Zittan et al, 2016 ²³	HMSA (commercial kit, Prometheus Laboratories)	No (30.9) ^a	—	—	Mucosa healing
Ungar et al ³⁸	—	—	—	—	Mucosa healing
Mazor et al ¹¹	(adapted) Antihuman lambda chain-based ELISA	No (30.5 samples) ^b no (12.7 samples) ^c	$\geq 3 \mu\text{g/mL}$ (b)	98/20.6	(a) Remission; (b) Active disease
Roblin et al ³⁶	Bridging ELISA (commercial kit)	9 (22.5)	No	No	(a) Absence of mucosa healing (b) Clinical remission
Imaeda et al, 2014 ²⁹	ELISA (+acid dissociation)	35 (23)	No	No	CRP $\leq 0.3 \text{ mg/dL}$

^aCutoff for ATA positivity >1 U/mL.
^bCutoff for ATA positivity >1.5 $\mu\text{g/mL}$ -eq.
^cCutoff for ATA positivity >3 $\mu\text{g/mL}$ -eq.




FIGURE 4. Methodologies for antidrug antibodies quantification.

significantly associated with a ≥ 70 -point increase in the mean total CD activity index score between infusions ($P < 0.001$).

In measurements performed at weeks 14 and/or 22, cutoff values varied from $<3^{26}$ to $\geq 7 \mu\text{g/mL}^{24}$ (Table 1). Patients with $\text{TL} > 3 \mu\text{g/mL}$ at weeks 14 and/or 22 had an approximately 66% lower likelihood to lose their response to IFX than those with subtherapeutic levels.²⁶ These findings are similar to data from the post-hoc analysis of the ACCENT I trial (A Crohn's Disease Clinical Trial Evaluating Infliximab in a New Long-term Treatment Regimen I),²⁷ which found that an IFX level $> 3.5 \mu\text{g/mL}$ at week 14 was a good predictor of sustained response at week 54. Patients with sustained response to scheduled maintenance IFX at 5 mg/kg had higher median IFX TL than those who lost response during the 54 weeks follow-up (4.0 versus 1.9 $\mu\text{g/mL}$, $P = 0.0331$). Adedokun et al²⁵ also measured IFX levels at weeks 30 and 54 and the levels, related with clinical response, were 3.7 and 1.7 $\mu\text{g/mL}$, respectively (Table 1). They suggested that more weight should be given to the threshold estimate at week 30 (3.7 $\mu\text{g/mL}$) because it was most representative of the steady-state trough concentration for both Active Ulcerative Colitis Trial studies. Therefore, patients with IFX TL $> 3.7 \mu\text{g/mL}$ at week 30 are more than twice as likely to have clinical response than patients with IFX TL $< 3.7 \mu\text{g/mL}$.

Mucosa Healing

Imaeda et al⁶ found that IFX TL $> 4 \mu\text{g/mL}$ was a good predictor of mucosa healing (Table 1). The authors also showed that the deterioration of the endoscopic findings was significantly associated with lower IFX TL. Two abstracts reported similar cutoff levels.^{33,38} In both of them, IFX levels were significantly higher in the mucosal healing group than in patients with active disease (4.3 versus 1.7 $\mu\text{g/mL}$, $P = 0.0002^{38}$).

Loss of Response

Steenholdt et al³⁵ showed that IFX TL < 0.5 in CD and < 0.8 in UC were good predictors of loss of response. IFX TL were significantly higher in both patients with CD and patients with UC who had maintained response to IFX compared with those who had lost response (median 2.8 $\mu\text{g/mL}$ versus median 0 $\mu\text{g/mL}$, for CD; and 3.8 $\mu\text{g/mL}$ versus 0 $\mu\text{g/mL}$ for UC). Higher levels were identified by Bortlik et al²⁶ ($< 3 \mu\text{g/mL}$) and Roblin et al³² ($< 2 \mu\text{g/mL}$).

Biomarkers

Imaeda et al⁶ showed that IFX levels $> 0.6 \mu\text{g/mL}$ could predict normalized CRP levels ($< 3 \text{ mg/dL}$) with good sensitivity and specificity (Table 1). C-reactive protein levels were significantly higher in the nonmucosal healing group than in the mucosal healing group (0.09 versus 1.32 mg/dL). Levesque et al³⁰ showed that IFX concentrations < 2.7 to $2.8 \mu\text{g/mL}$ predicted serum CRP levels $> 5 \text{ mg/L}$. Therefore, they suggested that a mean IFX trough concentration $< 3 \mu\text{g/mL}$ at week 8 was significantly associated with a higher probability for serum CRP concentrations $> 5 \text{ mg/L}$ at that time point. In a study by Vande Casteele et al,²² an IFX TL $> 2.79 \mu\text{g/mL}$ was considered to be a good predictor of CRP $< 5 \text{ mg/L}$, meaning that patients with IFX levels $< 2.79 \mu\text{g/mL}$ in a "current" sample were at higher risk of not achieving remission, defined as CRP $< 5 \text{ mg/L}$.

Adalimumab Levels

ADA information is sparse. Imaeda et al²⁹ evaluated 40 adults with CD and performed an ROC curve analysis to identify threshold levels of ADA that could predict normalized CRP levels (i.e., CRP $\leq 3 \text{ mg/dL}$). ADA levels $> 5.9 \mu\text{g/mL}$ predicted normalized CRP with high specificity (Table 2). Mazor et al¹¹ and

Roblin et al³⁶ conducted cross-sectional studies in patients taking ADA maintenance therapy; Mazor et al¹¹ enrolled patients with CD and Roblin et al³⁶ enrolled patients with CD or UC. In the study by Mazor et al, ADA TL >5.85 µg/mL predicted remission with a specificity and sensitivity of 70.6% and 68.0%, respectively. Roblin et al³⁶ showed that ADA serum concentrations <4.9 µg/mL predicted an absence of mucosa healing. The median ADA TL was significantly higher in cases of mucosa healing (6.5 versus 4.2 µg/mL in those without mucosa healing; $P < 0.005$). Moreover, serum levels higher than 4.85 µg/mL predicted clinical remission, defined as CD activity index <150 points or total Mayo score <3 (Table 2). Higher ADA TL were found in the work by Zittan et al (14.7 µg/mL in the mucosa healing group, versus 3.4 µg/mL in the non-MH group, $P = 6.25 \times 10^{-5}$).²³ Furthermore, Zittan et al suggested that ADA TL <8.14 µg/mL predicted MH with high sensitivity (Table 2). In the work by Ungar et al,³⁸ ADA levels >7.1 µg/mL identified patients with mucosa healing with 85% specificity. He also found that the association between higher levels of ADA and increased rate of mucosa healing reached a plateau at 12 µg/mL.

Incidence of ATIs and ATAs

Antidrug antibodies are described as the main cause of loss of response to biologic drugs over time. However, the incidence of antidrug antibodies varies significantly between studies. Taking into consideration those included in this systematic review, the ATI incidence varied from 9%²⁷ to 63.5%³⁷ (Table 1). In Bortlik et al,²⁶ 17% of the patients had ATIs but 55% were considered inconclusive. Marits et al⁸ reported 22 out of 28 patients with ATIs, wherein 18% of them were ATI transient, meaning that patients presented with ATIs in their serum which at some point disappeared. The same was reported by Vande Casteele et al,³⁴ where 15 of the 53 patients considered with ATIs were transient (Table 1). Vande Casteele et al²² reported an ATI incidence of 23.7%, and the authors were able to distinguish 4 groups of patients based on ATI and IFX status (Table 1). Paul et al³⁷ showed a global incidence of ATIs of 32.8%, considering a cutoff for ATIs of 10 ng/mL; with a cutoff of 5 ng/mL, the incidence was 63.5%. In the pediatric setting, 10% of the patients had ATIs in their blood at week 14, but the incidence increased to 16% at week 54.²⁴

Although ADA is a fully human antibody, some patients develop ATAs. Imaeda et al²⁹ described a 23% incidence of ATAs. Roblin et al³⁶ found a similar value (22.5%). In the cross-sectional study from Mazor et al,¹¹ 12.7% of the samples had ATA levels ≥3 µg/mL; when a cutoff of ≥1.5 µg/mL was established, the incidence rose to 30.5% (Table 2). Zittan et al²³ described an ATA incidence of 30.9%, using a cutoff of >1 U/mL.

Cutoff Levels of ATIs and ATAs

Only 5 studies^{11,22,34,35,37} performed an ROC curve analysis to identify threshold levels for antidrug antibodies. Steenholdt et al³⁵ reported that ATI levels, measured with fluid-phase RIA, >10 U/mL in patients with CD predicted “loss of clinical

response” with a specificity of 90% and sensitivity of 81%. In the subgroup of patients with UC, the specificity was higher (Table 1). ATI were significantly lower in both patients with CD and patients with UC who had maintained response to IFX compared with those who had lost response (median 0 U/mL versus median 35 U/mL for CD, and median 0 U/mL versus median 85 U/mL for UC).

Paul et al³⁷ also performed an ROC curve analysis using “loss of response” as the target clinical outcome. The authors suggested that ATI levels >200 ng/mL, assessed by the ELISA assay, predicted loss of response with a high specificity but with a low sensitivity (Table 1). A combined analysis was also performed on patients with CD with IFX levels <2 µg/mL and ATI levels <200 ng/mL. The ATIs predicted clinical remission with a high specificity and sensitivity (Table 1); patients with UC showed higher specificity (100%) but lower sensitivity (70%). The same analysis using “mucosa healing” as the clinical outcome was also supplied (Table 1). An ROC curve analysis for a threshold >9.1 U/mL at the time of loss of response predicted an “unsuccessful intervention” with a specificity of 82% and a sensitivity of 65%.³⁴ Therefore, patients having ATI TL >9.1 U/mL at the time of loss of response had a likelihood ratio of 3.6 for an unsuccessful intervention. It was also reported that patients with ATI levels <3.15 U/mL had a higher probability of being in remission.²¹

With regard to ADA, Mazor et al¹¹ suggested that a cutoff level ≥3 µg/mL, when using an adapted anti-human lambda chained-based ELISA assay, predicted active disease with high specificity but low sensitivity (Table 2). The authors showed a negative correlation between ADA drug levels and ATA levels and found that for patients with ATA levels ≥3 µg/mL-eq, the maximal ADA level was only 0.5 µg/mL.

DISCUSSION

The importance of measuring drug levels and antidrug antibody levels to adjust therapy is undisputable. The major hindrance to its implementation in daily clinical practice is the lack of a universally valid assay and the absence of a cutoff level clearly related with a clinical outcome. One cannot easily compare results from different studies, as they use distinct assays that have different limitations and lower limits of quantification.

Regarding the measurement of IFX levels, classic ELISA is the methodology most frequently used, but other methods are available, such as HMSA and fluid-phase RIA. Studies^{40–42} that have compared performance of different methods to measure drug levels have concluded the same; there is a good qualitative correlation between different assays (e.g., IFX detection rates of 76% with ELISA and 82% with RIA⁴²). Furthermore, in some cases, there is a good quantitative correlation (e.g., ELISA and RIA, $R^2 = 0.98$, $P = 0.001$ ⁴⁰; ELISA and RIA, Pearson $r = 0.91$, $P < 0.0001$ ⁴¹) but not a perfect agreement on drug concentrations (e.g., maximum difference of 1.41 µg/mL between ELISA and RIA⁴⁰), and this emphasizes the importance of

establishing different cutoff levels according to the methodology used. The threshold levels assessed by ROC curve analysis were quite different between the studies. This can be due to (1) different methodology (even using the same principle, such as bridging ELISA, home-made ELISA, and commercial kits), (2) different study design and sample characteristics, and/or (3) different endpoints. This heterogeneity justifies the obstacle to perform a meta-analysis. A systematic review and meta-analysis was recently published on this topic and suggested a cutoff level of 2 $\mu\text{g/mL}$ to predict remission ($\text{RR} = 2.9$, 95% confidence interval, 1.8–4.7, $P < 0.001$), but there was a high statistical heterogeneity ($I^2 = 88\%$).⁴³ However, TL were always associated with a better clinical endpoint: clinical remission, mucosa healing, normalized CRP, or loss of response. Our review emphasizes the importance of measuring drug levels during maintenance therapy as well as in cases of loss of response, cases with persistent high levels of CRP, and when mucosal lesions are still present. In the induction phase, the only study reported did not show any advantage of measuring IFX at 2 weeks because this corresponded to the loading period and it was not possible to differentiate responders from nonresponders. However, at weeks 8, 14, and 30, the different studies found significant differences between responders and nonresponders, and one of these time periods should be chosen by clinicians for strategic therapeutic decisions, namely increasing drug dose or addition of 1 immunomodulatory drug. Active Ulcerative Colitis Trial subanalysis suggested week 30 is ideal and argued that this time corresponds to the steady state of the drug. Two studies (TAILORIX⁴⁴ and TAXIT⁴⁵) have concluded that in maintenance phase, concentration-based dose adjustment was not superior to dose adjustment based on symptoms alone. However, TAXIT trial also showed that patients in the “clinically based dosing” group had more flares during the course of treatment than those in the “concentration-based dosing” group.

Overall, there is evidence for determining drug levels in weeks 6, 14, 22, 30, and 54. During maintenance, therapeutic drug monitoring should be considered in case of loss of response, mucosal ulceration, and elevated biomarkers, such as CRP and fecal calprotectin (Fig. 5).

Figure 6 shows how therapeutic drug monitoring may be used to highlight factors influencing loss of response. Two branches are schematized: for patients with loss of response and high levels of drug (pink branche) and for patients with loss of response and low drug levels (green branche). Pharmacodynamic, pharmacokinetic, and immunogenicity factors may be identified and help clinicians to handle therapeutic decisions.

All methodologies available (ELISA, HMSA, fluid-phase RIA) seem qualitatively equivalent, so either one can be used to monitor drug levels. However, the clinician should take into consideration that there are disagreements on IFX concentration between assays, therefore for each patient, drug levels should be always measured with the same assay. Concerning antidrug antibody levels, the variability among methods is more significant. Enzyme-linked immunosorbent assay (ELISA) is the methodology most frequently used; however, not all ELISAs use the same principle. A bridging ELISA, or double antigen ELISA, uses the drug, in this case, IFX or ADA as the captured antigen and as the detection antibody. Consequently, this method is susceptible to several limitations, namely false-positive results, caused by rheumatoid factors or activated complement fragments that cross-bind the drug's fragment crystallizable region. False-negative results are due to the assay's inability to detect monovalent immunoglobulin G4 (IgG4) and antidrug antibodies in the presence of the drug. This method was used by 6 of the studies included in this systematic review.^{24–27,36,37} Since this assay has no sensitivity to detect antidrug antibodies in the presence of the drug, some of the studies did not measure antidrug antibodies if there were drug levels in the serum and considered those samples




FIGURE 5. Time points for drug level determination.




FIGURE 6. Therapeutic drug monitoring brought to light the factors influencing the loss of response.

as "ATI inconclusive." Cornillie et al²⁷ considered samples that had IFX levels $>0.1 \mu\text{g/mL}$ as "ATI inconclusive," whereas Bortlik et al²⁶ only considered samples that had IFX levels $>3 \mu\text{g/mL}$ as "ATI inconclusive." Given that half of the patients in clinical trials had the drug in their serum, the use of a bridging ELISA for anti-IFX detection may lead to serious bias. This must be taken into consideration when one tries to draw conclusions about the therapeutic importance of ATIs using bridging methodology.³⁵

We should also keep in mind that study populations and study designs were different. Some included only patients with CD or UC, whereas others comprised both types of patients; some were prospective cohorts while others were cross-sectional studies or post-hoc analyses of controlled trials. These differences can explain why the incidence of antidrug antibodies was so varied between them, even when using the same assay. For example, both studies from Cornillie et al²⁷ and Paul et al³⁷ used a bridging ELISA to measure antidrug antibodies but the incidence of ATI positivity was 9% and 32.8%, respectively. This could be explained by the fact that the first study was a post-hoc analysis

of the ACCENT I trial that enrolled 573 adult patients with CD, whereas the second was a prospective cohort with 103 adults with CD or UC.

Kopylov et al⁴⁶ developed a different ELISA method, anti-human lambda chain-based ELISA, to overcome the false-negative results associated with the presence of the drug. The authors took advantage of the fact that antidrug antibodies have a lambda light chain, whereas the drug has a kappa light chain, and they used an anti-lambda antibody as the detection antibody, ensuring that they were only measuring antidrug antibodies. Mazar et al¹¹ adapted this method to measure antibodies to ADA. Those authors described an incidence for ATA positivity of 30.5%, which showed the sensitivity of anti-lambda chain ELISA and its low rate of drug interference. However, in serum with high levels of a drug, even anti-lambda chain ELISA is not able to completely overcome drug interference.⁴⁷ Anti-lambda chain ELISA is also unable to detect anti-idiotypic antibodies, i.e., antibodies that recognize functional binding epitopes.⁴⁷

One way of overcoming drug interference is to perform a prior acidic dissociation. Imaeda et al pretreated samples with

acid in both the IFX study and ADA study.^{6,29} In a previous work, the authors showed the ability of this new method to detect ATIs in samples containing detectable levels of IFX, which proved to be more accurate than the bridging ELISA.⁴⁸ From a total of 58 samples, the methodology by Imaeda et al could detect an additional 14 positive samples, of which, by the bridging ELISA, 8 had been considered negative and 6 “inconclusive.”

Three studies^{24,30,34} used the HMSA to measure ATIs, an alternative assay to ELISA. The HMSA uses size exclusion high-performance liquid chromatography.⁴⁷ Although HMSA requires expensive equipment, the authors of those studies state many advantages, including the ability to overcome many potential artifacts encountered in the solid-phase ELISA, the ability to detect high and low affinity antibodies (low affinity antibodies may not be detected by ELISA due to multiple washing steps), the detection of all immunoglobulin isotypes and all IgG subclasses (including IgG4), and the fact that it is not affected by substances present in serum.⁴⁹ However, a different ATI incidence was reported by the 3 studies, which can be explained by differences in the study population and sample size (Table 1).

Another assay is able to bridge the gaps of the ELISA methodology. In fluid-phase radioimmunoassay (RIA), used by Steenholdt et al,³⁵ a radio-labeled antibody to detect and quantify the amount of antidrug antibodies is applied. It has proved to be more sensitive than ELISA, as it is able to detect antidrug antibodies in the presence of the drug and IgG4 isotype.^{40,41} Moreover, fluid-phase RIA overcomes matrix effects encountered in solid-phase assays due to epitope masking via protein aggregation. The major limitation of RIA is the need for advanced laboratory facilities.⁴⁷

Therefore, the differences in methodology, study design, and sample size and characteristics may also explain why the 4 studies with IFX^{22,34,35,37} that performed an ROC curve analysis in order to find a cutoff level of antidrug antibodies related with a clinical outcome found different threshold levels. It is also not easy to compare the thresholds between studies because they used different units (U/mL; μ g/mL; ng/mL) and defined different endpoints. A serious limitation of all of the studies was the inability to show whether or not antidrug antibodies were neutralizing.

It is important to address whether or not antidrug antibodies are functional, because we know that antidrug antibody detection in serum does not always correlate with loss of clinical response.^{47,50} Moreover, sometimes the presence of antidrug antibodies may actually increase the half-life of the drug; if 1 or 2, but not more, antidrug antibodies bind to the drug, the complex will bind to Neonatal fragment crystallizable receptor and will escape elimination.⁵¹ A study comparing different methodologies (ELISA, EIA, RGA, RIA) to measure antidrug antibody levels has been published and concluded that the ability to detect anti-ATIs is comparable with respect to basic analytical properties. ELISA and RIA showed a good correlation ($R^2 = 0.73$, $P = 0.03$), but the agreement was not so good, with a mean titer difference of -2400 (-5000 to 200), which can be partially explained by the inability of bridging ELISA to detect IgG4

antidrug antibodies. The authors suggest that clinicians should choose an assay where assessments take place in fluid phase and where all anti-IFX IgG isotypes are quantified.⁴⁰

CONCLUSION

Currently, there is no doubt that drug levels correlate with clinical and endoscopic outcomes, and this knowledge is the basis of drug monitoring. Nevertheless, it can only be widely used in clinical practice when there is a consensus on the thresholds of drug and antidrug antibody levels that correlate with a specific clinical outcome, including either clinical remission or loss of response. Concerning drug level monitoring, any methodology is adequate but the data published by now is insufficient to come up with a cutoff level. With respect to antidrug antibody levels, assays have significantly different sensitivity, therefore it will be necessary to define a gold standard method or to establish different cutoff levels for different methodologies.

REFERENCES

- Colombel JF, Sandborn WJ, Reinisch W, et al. Infliximab, azathioprine, or combination therapy for Crohn's disease. *N Engl J Med*. 2010;362:1383–1395.
- Hanauer SB, Feagan BG, Lichtenstein GR, et al. Maintenance infliximab for Crohn's disease: the ACCENT I randomised trial. *Lancet*. 2002;359:1541–1549.
- Rutgeerts P, Sandborn WJ, Feagan BG, et al. Infliximab for induction and maintenance therapy for ulcerative colitis. *N Engl J Med*. 2005;353:2462–2476.
- Peters CP, Eshuis EJ, Toxopeus FM, et al. Adalimumab for Crohn's disease: long-term sustained benefit in a population-based cohort of 438 patients. *J Crohns Colitis*. 2014;8:866–875.
- García-Bosch O, Gisbert JP, Cañas-Ventura A, et al. Observational study on the efficacy of adalimumab for the treatment of ulcerative colitis and predictors of outcome. *J Crohns Colitis*. 2013;7:717–722.
- Imaeda H, Bamba S, Takahashi K, et al. Relationship between serum infliximab trough levels and endoscopic activities in patients with Crohn's disease under scheduled maintenance treatment. *J Gastroenterol*. 2014;49:674–682.
- Seow CH, Newman A, Irwin SP, et al. Trough serum infliximab: a predictive factor of clinical outcome for infliximab treatment in acute ulcerative colitis. *Gut*. 2010;59:49–54.
- Marits P, Landucci L, Sundin U, et al. Trough s-infliximab and antibodies towards infliximab in a cohort of 79 IBD patients with maintenance infliximab treatment. *J Crohns Colitis*. 2014;8:881–889.
- Nanda KS, Cheifetz AS, Moss AC. Impact of antibodies to infliximab on clinical outcomes and serum infliximab levels in patients with inflammatory bowel disease (IBD): a meta-analysis. *Am J Gastroenterol*. 2013;108:40–47.
- Ben-Horin S, Yavzori M, Katz L, et al. The immunogenic part of infliximab is the F(ab')₂, but measuring antibodies to the intact infliximab molecule is more clinically useful. *Gut*. 2011;60:41–48.
- Mazor Y, Almog R, Kopylov U, et al. Adalimumab drug and antibody levels as predictors of clinical and laboratory response in patients with Crohn's disease. *Aliment Pharmacol Ther*. 2014;40:620–628.
- Felice C, Marzo M, Pugliese D, et al. Therapeutic drug monitoring of anti-TNF- α agents in inflammatory bowel diseases. *Expert Opin Biol Ther*. 2015;15:1107–1117.
- Yarur AJ, Abreu MT, Deshpande AR, et al. Therapeutic drug monitoring in patients with inflammatory bowel disease. *World J Gastroenterol*. 2014;20:3475–3484.
- Ben-Horin S, Chowers Y. Tailoring anti-TNF therapy in IBD: drug levels and disease activity. *Nat Rev Gastroenterol Hepatol*. 2014;11:243–255.
- Ben-Horin S. Loss of response to anti-tumor necrosis factors: what is the next step? *Dig Dis*. 2014;32:384–388.

16. Bendtzen K. Immunogenicity of anti-TNF- α Biotherapies: I. Individualized Medicine based on Immunopharmacological evidence. *Front Immunol.* 2015;6:152.
17. Khanna R, Sattin BD, Afif W, et al. Review article: a clinician's guide for therapeutic drug monitoring of infliximab in inflammatory bowel disease. *Aliment Pharmacol Ther.* 2013;38:447–459.
18. Pallagi-Kunstár É, Farkas K, Szepes Z, et al. Utility of serum TNF- α , infliximab trough level, and antibody titers in inflammatory bowel disease. *World J Gastroenterol.* 2014;20:5031–5035.
19. Vande Casteele N, Feagan B, Gils A, et al. Therapeutic drug monitoring in inflammatory bowel disease: current state and future perspectives. *Curr Gastroenterol Rep.* 2014;16:1–8.
20. Altwegg R, Vincent T. TNF blocking therapies and immunomonitoring in patients with inflammatory bowel disease. *Mediators Inflamm.* 2014;2014:172821.
21. Stroup D, Berlin J, Morton S, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis of Observational Studies in Epidemiology (MOOSE) group. *JAMA.* 2000;283:2008–2012.
22. Vande Casteele N, Khanna R, Levesque BG, et al. The relationship between infliximab concentrations, antibodies to infliximab and disease activity in Crohn's disease. *Gut.* 2015;64:1539–1545.
23. Zittan E, Kabakchiev B, Milgrom R, et al. Higher adalimumab drug levels are associated with mucosal healing in patients with Crohn's disease. *J Crohns Colitis.* 2016;10:510–515.
24. Singh N, Rosenthal CJ, Melmed GY, et al. Early infliximab trough levels are associated with persistent remission in pediatric patients with inflammatory bowel disease. *Inflamm Bowel Dis.* 2014;20:1708–1713.
25. Adedokun OJ, Sandborn WJ, Feagan BG, et al. Association between serum concentration of infliximab and efficacy in adult patients with ulcerative colitis. *Gastroenterology.* 2014;147:1296–1307.e5.
26. Bortlik M, Duricova D, Malickova K, et al. Infliximab trough levels may predict sustained response to infliximab in patients with Crohn's disease. *J Crohns Colitis.* 2013;7:736–743.
27. Cornillie F, Hanauer SB, Diamond RH, et al. Postinduction serum infliximab trough level and decrease of C-reactive protein level are associated with durable sustained response to infliximab: a retrospective analysis of the ACCENT I trial. *Gut.* 2014;63:1721–1727.
28. Echarri A, Ferreira R, Fraga R, et al. Impact of postinduction infliximab trough level and disease activity on primary response in Crohn's Disease. *J Crohns Colitis.* 2015;9:S342–S343.
29. Imaeda H, Takahashi K, Fujimoto T, et al. Clinical utility of newly developed immunoassays for serum concentrations of adalimumab and anti-adalimumab antibodies in patients with Crohn's disease. *J Gastroenterol.* 2014;49:100–109.
30. Levesque BG, Greenberg GR, Zou G, et al. A prospective cohort study to determine the relationship between serum infliximab concentration and efficacy in patients with luminal Crohn's disease. *Aliment Pharmacol Ther.* 2014;39:1126–1135.
31. Papamichail K, Vande Casteele N, Billiet T, et al. Early therapeutic drug monitoring for prediction of short-term mucosal healing in patients with ulcerative colitis treated with infliximab. *Gastroenterology.* 2015;148:S848.
32. Roblin X, Duru G, Clavel L, et al. Faecal calprotectin measurement and infliximab trough levels predict therapeutic evolution CD patients in clinical remission. *Drug Levels & Biomarkers.* 2015;148:S–853.
33. Tang J, Gao X, Zhi M, et al. Serum infliximab levels and early mucosal healing in Crohn's disease. *J Crohns Colitis.* 2014;8:S209–S210.
34. Vande Casteele N, Gils A, Singh S, et al. Antibody response to infliximab and its impact on pharmacokinetics can be transient. *Am J Gastroenterol.* 2013;108:962–971.
35. Steenholdt C, Bendtzen K, Brynskov J, et al. Cut-off levels and diagnostic accuracy of infliximab trough levels and anti-infliximab antibodies in Crohn's disease. *Scand J Gastroenterol.* 2011;46:310–318.
36. Roblin X, Marotte H, Rinaudo M, et al. Association between pharmacokinetics of adalimumab and mucosal healing in patients with inflammatory bowel diseases. *Clin Gastroenterol Hepatol.* 2014;12:80–84.e2.
37. Paul S, Del Tedesco E, Marotte H, et al. Therapeutic drug monitoring of infliximab and mucosal healing in inflammatory bowel disease: a prospective study. *Inflamm Bowel Dis.* 2013;19:2568–2576.
38. Ungar B, Levy I, Yavne Y, et al. Optimizing anti-TNF α therapy: serum levels of infliximab and adalimumab associate with mucosal healing in patients with inflammatory bowel diseases. *Clin Gastroenterol Hepatol.* 2016;14:550–557.
39. Chiu YL, Rubin DT, Vermeire S, et al. Serum adalimumab concentration and clinical remission in patients with Crohn's disease. *Inflamm Bowel Dis.* 2013;19:1112–1122.
40. Steenholdt C, Ainsworth MA, Tovey M, et al. Comparison of techniques for monitoring infliximab and antibodies against infliximab in Crohn's Disease. *Ther Drug Monit.* 2013;35:530–538.
41. Vande Casteele N, Buurman DJ, Sturkenboom MGG, et al. Detection of infliximab levels and anti-infliximab antibodies: a comparison of three different assays. *Aliment Pharmacol Ther.* 2012;36:765–771.
42. Steenholdt C, Bendtzen K, Brynskov J, et al. Clinical implications of measuring drug and anti-drug antibodies by different assays when optimizing infliximab treatment failure in Crohn's disease. *J Crohns Colitis.* 2014;8:S291.
43. Moore C, Corbett G, Moss AC. Systematic review and meta-analysis: serum infliximab levels during maintenance therapy and outcomes in inflammatory bowel disease. *J Crohns Colitis.* 2016;10:619–625.
44. D'Haens G, Vermeire S, Lambrecht G, et al. OP029 Drug-concentration versus symptom-driven dose adaptation of infliximab in patients with active Crohn's disease: a prospective, randomised, multicentre trial (Tailorix). *J Crohns Colitis.* 2016;10:24.
45. Casteele NV, Gils A, Ballet V, et al. OP001 Randomised controlled trial of drug level versus clinically based dosing of infliximab maintenance therapy in IBD: final results of the TAXIT study. *United Eur Gastroenterol J.* 2013:A1–A134.
46. Kopylov U, Mazor Y, Yavzori M, et al. Clinical utility of antihuman lambda chain-based enzyme-linked immunosorbent assay (ELISA) versus double antigen ELISA for the detection of anti-infliximab antibodies. *Inflamm Bowel Dis.* 2012;18:1628–1633.
47. Bendtzen K. Personalized medicine: theranostics (therapeutics diagnostics) essential rational use tumor necrosis factor-alpha antagonists. *Discov Med.* 2013;15:201–211.
48. Imaeda H, Andoh A, Fujiyama Y. Development of a new immunoassay for the accurate determination of anti-infliximab antibodies in inflammatory bowel disease. *J Gastroenterol.* 2012;47:136–143.
49. Wang SL, Ohrmund L, Hauenstein S, et al. Development and validation of a homogeneous mobility shift assay for the measurement of infliximab and antibodies-to-infliximab levels in patient serum. *J Immunol Methods.* 2012;382:177–188.
50. Ungar B, Chowes Y, Yavzori M, et al. The temporal evolution of anti-drug antibodies in patients with inflammatory bowel disease treated with infliximab. *Gut.* 2014;63:1258–1264.
51. Wang W, Wang EQ, Balthasar JP. Monoclonal antibody pharmacokinetics and pharmacodynamics. *Clin Pharmacol Ther.* 2008;84:548–558.

Inflammatory Bowel Diseases

Instructions for Authors

SCOPE

The focus of the journal *Inflammatory Bowel Diseases*® is to bring the most current information in the clinical and basic sciences to all physicians caring for patients with inflammatory bowel diseases, and to investigators performing research in inflammatory bowel diseases and related fields. The journal publishes peer-reviewed manuscripts and review articles in basic and clinical sciences, updates on clinical trials, reviews of the current literature, editorials, and other features.

ETHICAL/LEGAL CONSIDERATIONS

A submitted manuscript must be an original contribution not previously published (except as an abstract or preliminary report), must not be under consideration for publication elsewhere, and, if accepted, must not be published elsewhere in similar form, in any language, without the consent of the publisher. Each person listed as an author is expected to have participated in the study to a significant extent. Although the editors and referees make every effort to ensure the validity of published manuscripts, the final responsibility rests with the authors, not with the journal, its editors, or the publisher. All manuscripts must be submitted through the journal's online submission system at <http://www.editorialmanager.com/ibd/>.

Inflammatory Bowel Diseases utilizes the plagiarism software, CrossCheck. Manuscripts submitted to this journal may be screened for plagiarism against previously published works. Authors who have been found to be guilty of plagiarism after the appropriate institutional investigation will be banned from submitting to the journal for a defined period of time.

Patient Anonymity and Informed Consent - It is the author's responsibility to ensure that a patient's anonymity be carefully protected and to verify that any experimental investigation with human subjects reported in the manuscript was performed with informed consent and following all the guidelines for experimental investigation with human subjects required by the institution(s) with which all the authors are affiliated. Authors should mask patients' eyes and remove patients' names from figures unless they obtain written consent from the patients and submit written consent with the manuscript.

Human Studies - Human experimentation must conform to ethical standards, and be approved by the appropriate Institutional Review Board (IRB). A statement concerning IRB approval and consent procedures must appear at the beginning of the Methods section. Any systematic data gathering effort in patients or volunteers must be approved by an IRB or adhere to appropriate local/national regulations. The Editors of *Inflammatory Bowel Diseases* are concerned about appropriate IRB review and informed consent. Authors may be questioned about the details of consent forms or the consent process. On occasion, the Editor-in-Chief may request a copy of the approved IRB application from the author. Lack of appropriate consent or documentation may be grounds for rejection. Local IRB approval does not guarantee acceptability; the final decision will be made by the Editor-in-Chief.

Animal Studies - Experimental work on animals must conform to the guidelines laid out in the Guide for the Care and Use of Laboratory Animals, which is available from the National

Academy of Science; a text-only version is available at <http://www.nap.edu/readingroom/books/labrats/>. Adherence to all relevant regulations and/or approval of the appropriate institutional Animal Care Committee or governmental licensure of the investigator and/or laboratory must be obtained. A statement concerning such approval must be included at the beginning of the Methods section. The Editors of *Inflammatory Bowel Diseases* are concerned about appropriate animal care. On occasion, the Editor-in-Chief may request a copy of the approved Animal Care Committee application from the author. Local committee approval does not guarantee acceptability; the final decision will be made by the Editor-in-Chief.

Registration of Clinical Trials - All clinical trials that involve investigational drugs supported by a pharmaceutical firm or investigational devices supported by a device manufacturer must be registered at the time that a manuscript is submitted to *Inflammatory Bowel Diseases* for publication. The registry and registration number must be stated in the first paragraph of the Methods section of the manuscript.

Compliance with NIH and Other Research Funding Agency Accessibility

Requirements - A number of research funding agencies now require or request authors to submit the post-print (the article after peer review and acceptance but not the final published article) to a repository that is accessible online by all without charge. As a service to our authors, LWW will identify to the National Library of Medicine (NLM) articles that require deposit and will transmit the post-print of an article based on research funded in whole or in part by the National Institutes of Health, Wellcome Trust, Howard Hughes Medical Institute, or other funding agencies to PubMed Central. The revised Copyright Transfer Agreement & Disclosure form provides the mechanism.

Copyright - All authors must complete and submit the journal's copyright transfer agreement, which includes a section on the disclosure of potential conflicts of interest based on the recommendations of the International Committee of Medical Journal Editors, "Uniform Requirements for Manuscripts Submitted to Biomedical Journals" (www.icmje.org/update.html).

A copy of the form is made available to the submitting author within the Editorial Manager submission process. Co-authors will automatically receive an Email with instructions on completing the form upon submission.

Conflicts of Interest - Authors must state all possible conflicts of interest in the manuscript, including financial, consultant, institutional and other relationships that might lead to bias or a conflict of interest. If there is no conflict of interest, this should also be explicitly stated as none declared. All sources of funding should be acknowledged in the manuscript. All relevant conflicts of interest and sources of funding should be included on the title page of the manuscript with the heading "Conflicts of Interest and Source of Funding".

For example: Conflicts of Interest and Source of Funding: A has received honoraria from Company Z. B is currently receiving a grant (#12345) from Organization Y, and is on the speaker's bureau for Organization X — the CME organizers for Company A. For the remaining authors none were declared.

In addition, each author must complete and submit the journal's copyright transfer agreement, which includes a section on the disclosure of potential conflicts of interest based on the recommendations of the International Committee of Medical Journal Editors, "Uniform

Requirements for Manuscripts Submitted to Biomedical Journals"
(www.icmje.org/update.html).

A copy of the form is made available to the submitting author within the Editorial Manager submission process. Co-authors will automatically receive an Email with instructions on completing the form upon submission.

Permissions - Authors must submit written permission from the copyright owner (usually the publisher) to use direct quotations, tables, or illustrations that have appeared in copyrighted form elsewhere, along with complete details about the source. Any permissions fees that might be required by the copyright owner are the responsibility of the authors requesting use of the borrowed material, and not the responsibility of the publisher.

PREPARATION OF MANUSCRIPT

Manuscripts that do not adhere to the following instructions will be returned to the corresponding author for technical revision before undergoing peer review. See specific guidelines for Letters to the Editor below. For questions on preparing manuscripts for submission, please contact Meghann Knowles at meghann@jjeditorial.com.

Title Page Include on the title page (a) complete manuscript title; (b) authors' full names, highest academic degrees, and affiliations; (c) name and address for correspondence, including fax number, telephone number, and e-mail address; and (d) sources of support that require acknowledgment.

The title page must also include disclosure of funding received for this work from any of the following organizations: National Institutes of Health (NIH); Wellcome Trust; Howard Hughes Medical Institute (HHMI); and other(s).

Structured Abstract and Key Words Limit the abstract to 250 words. It must be factual and comprehensive. Do not cite references in the abstract. Limit the use of abbreviations and acronyms, and avoid general statements (e.g., "the significance of the results is discussed"). It should be sectioned into Background, Methods, Results, and Conclusions. List three to five key words.

Text For full-length research articles, organize the manuscript in the following sequence: Abstract and Key Words, Introduction, Materials and Methods, Ethical Considerations, Results, Discussion, Acknowledgment, References, Tables, and Figure Legends. Original Research Articles: Authors are encouraged to submit articles in basic or clinical science. Review Articles of exceptional merit will also be accepted. Contributors should communicate with the Editors before submitting a review. Abbreviations must be defined at first mention in text and in each table and figure. If a brand name is cited, manufacturer and address (city and state/country) should be supplied. Acknowledge all forms of support, including pharmaceutical industry support, in an Acknowledgment paragraph.

Abbreviations For a list of standard abbreviations, consult the Council of Biology Editors Style Guide (available from the Council of Science Editors, 9650 Rockville Pike, Bethesda, MD 20814) or other standard sources. Write out the full term for each abbreviation at its first use unless it is a standard unit of measure.

References The authors are responsible for the accuracy of the references. Key the references (double-spaced) at the end of the manuscript. Cite the references in text in the order of appearance, including those references cited in tables and figure legends at the chronological citation of the tables and figures in text. Cite unpublished data, such as papers submitted but not yet accepted for publication, personal communications, in parentheses in the text. If there are more than three authors, only name the first three authors and then use et al. For abbreviations of journal names, refer to List of Journals Indexed in Index Medicus. This can be accessed at <http://www.nlm.nih.gov/tsd/serials/lji.html>.

Sample references are given below:

Journal Article

1. Gudlaugsdottir S, van Dekken H, Stijnen T, et al. Prolonged use of proton pump inhibitors, CagA status, and the outcome of *Helicobacter pylori* gastritis. *J Clin Gastroenterol*. 2002;34:536-540.

Book Chapter

2. Tobin RW, Kimmey MB. Painful diseases of the gastrointestinal tract. In: Loeser JD, ed. *Bonica's Management of Pain*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2001:1269-1292.

Entire Book

3. Rohen JW, Yokochi C, Lütjen-Drecoll E. *Color Atlas of Anatomy: A Photographic Study of the Human Body*. 5th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2002.

Software

4. Epi Info [computer program]. Version 6. Atlanta: Centers for Disease Control and Prevention; 1994.

Online Journals

5. Friedman SA. Preeclampsia: a review of the role of prostaglandins. *ObstetGynecol* [serial online]. January 1988;71:22-37. Available from: BRS Information Technologies, McLean, VA. Accessed December 15, 1990.

Database

6. CANCERNET-PDQ [database online]. Bethesda, MD: National Cancer Institute; 2014. Updated March 29, 2014.

World Wide Web

7. Gostin LO. Drug use and HIV/AIDS [JAMA HIV/AIDS Web site]. June 1, 2015. Available at: <http://www.ama-assn.org/special/hiv/ethics>. Accessed July 26, 2015.

Figures Art should be created/scanned and saved and submitted as either a TIFF (tagged image file format), or an EPS (encapsulated postscript) file. Line art must have a resolution of at least 1200 dpi (dots per inch), and electronic photographs, radiographs, CT scans, and other scanned images must have a resolution of at least 300 dpi. If fonts are used in the artwork, they must be converted to paths or outlines or they must be embedded in the files. Color images must be created/scanned and saved and submitted as CMYK files. Please note that artwork generated from office suite programs such as Corel Draw and MS Word and artwork downloaded from the Internet (JPEG or GIF files) cannot be used. Cite figures consecutively in the manuscript, and number them in the order in which they are discussed.

Figure Legends Legends must be submitted for all figures. They should be brief and specific, and they should appear after the tables. Use scale markers in the image for electron micrographs, and indicate the type of stain used.

Color Figures At no cost to the author, the journal accepts for publication color figures that will enhance the article. Figures are also published online in color at no cost.

Steps for Submitting Artwork

1. Please reference "5 Steps to Creating Digital Artwork (pdf)" available at <http://edmgr.ovid.com/lww-final/accounts/5StepsforArt.pdf>, specifically steps 2 and 3, when creating figures for submission.
2. Create, Scan and Save according to the "5 Steps to Creating Digital Artwork (pdf)".
3. Compare your final figure to the Target Digital-Imaging Results listed below.
4. Upload each figure to the journal's online submission system in conjunction with your manuscript text and tables.

Digital Artwork Guideline Checklist

Here are the basics to have in place before submitting your digital art to *Inflammatory Bowel Diseases*. Artwork saved as TIFF and EPS files. Do not save TIFFs as compressed files. PowerPoint files are also acceptable.

- Artwork created as the actual size (or slightly larger) it will appear in the journal. (To get an idea of the size images should be when they print, study a copy of the journal to which you wish to submit. Measure the artwork typically shown and scale your image to match.)
- Crop out any white or black space surrounding the image.
- Text and fonts in any figure are one of the acceptable fonts: Helvetica, Times Roman, Symbol, Mathematical PI, and European PI.
- Color images are created/scanned and saved and submitted as CMYK only. Do not submit any figures in RGB mode because RGB is the color mode used for screens/monitors and CMYK is the color mode used for print.
- Line art saved at a resolution of at least 1200 dpi.
- Images saved at a resolution of at least 300 dpi.
- Each figure saved as a separate file and saved separately from the accompanying text file.
- For multi-panel or composite figures only: Any figure with multiple parts should be sent as one file with each part labeled the way it is to appear in print.

Remember:

- Artwork generated from office suite programs such as CorelDRAW, MS Word, Excel, and artwork downloaded from the Internet (JPEG or GIF files) cannot be used because the quality is poor when printed.
- Cite figures consecutively in your manuscript.

- Number figures in the figure legend in the order in which they are discussed.
- Upload figures consecutively to the journal's online submission system and number figures consecutively in the Description box during upload.

Tables Create tables using the table creating and editing feature of your word processing software (e.g., Word). Do not use Excel or comparable spreadsheet programs. Do not submit tables as image files or images placed in Word documents, tables must be provided as editable text (Word files are preferred). Submit all tables as separate files. Cite tables consecutively in the text, and number them in that order. Key each on a separate sheet, and include the table title, appropriate column heads, and explanatory legends (including definitions of any abbreviations used). Do not imbed tables within the body of the manuscript. They should be self-explanatory and should supplement, rather than duplicate, the material in the text.

Supplemental Digital Content (SDC) Authors may submit supplementary materials via the journal's online submission system that enhance their article's text to be considered for online posting. SDC may include standard media such as text documents, graphs, audio, video, etc. On the Attach Files page of the submission process, please select Supplemental Audio, Video, or Data for your uploaded file as the Submission Item. If an article with SDC is accepted, our production staff will create a URL with the SDC file. The URL will be placed in the call-out within the article. SDC files are not copy-edited by LWW staff, they will be presented digitally as submitted. Please supply the SDC files as you would like them to appear in final publication (include legends in the same file as the images; make text double spaced or single spaced per your preference). For a list of all available file types and detailed instructions, please visit <http://links.lww.com/A142>.

SDC Call-outs Supplemental Digital Content must be cited consecutively in the text of the submitted manuscript. Citations should include the type of material submitted (Audio, Figure, Table, etc.), be clearly labeled as "Supplemental Digital Content," include the sequential list number, and provide a description of the supplemental content. All descriptive text should be included in the call-out as it will not appear elsewhere in the article.

Example: We performed many tests on the degrees of flexibility in the elbow (see Video, Supplemental Digital Content 1, which demonstrates elbow flexibility) and found our results inconclusive.

List of Supplemental Digital Content A listing of Supplemental Digital Content must be submitted at the end of the manuscript file. Include the SDC number and file type of the Supplemental Digital Content. This text will be removed by our production staff and not be published.

Example:

Supplemental Digital Content 1.wmv

SDC Files Requirements All acceptable file types are permissible up to 10 MBs. For audio or video files greater than 10 MBs, authors should first query the journal office for approval. For a list of all available file types and detailed instructions, please visit <http://links.lww.com/A142>.

Style Follow American Medical Association Manual of Style (10th edition). Stedman's Medical Dictionary (27th edition) and Merriam Webster's Collegiate Dictionary (10th edition) should be used as standard references. Refer to drugs and therapeutic agents by their accepted

generic or chemical names, and do not abbreviate them. Use code numbers only when a generic name is not yet available. In that case, supply the chemical name and a figure giving the chemical structure of the drug. Capitalize the trade names of drugs and place them in parentheses after the generic names. To comply with trademark law, include the name and location (city and state in USA; city and country outside USA) of the manufacturer of any drugs, supplies, or equipment mentioned in the manuscript. Use the metric system to express units of measure and degrees Celsius to express temperatures, and use SI units rather than conventional units.

Basic and Clinical IBD Review Articles Guidelines Basic and clinical IBD Review articles should present recent advances in a relatively narrow topic that have been made in cutting edge research.

The goal of the basic and clinical IBD review articles should be to present a complete summary of important research areas that are now improving our understanding of Crohn's disease and ulcerative colitis.

The body of basic and clinical IBD Review articles should be no longer than 20 double spaced pages (not including references, figures, and tables). There should be no more than 6 tables and figures (combined). Supplemental figures and tables will be allowed online. If page limits need to be increased, the authors may request permission from the Editors to increase the length of the Review article. The number of references should be limited to 100. The Review article should be focused on a single specific topic. All review articles will be peer-reviewed.

Future Directions and Methods for IBD Research Guidelines Original research articles and review articles on future directions and methods for IBD research should discuss important basic and clinical areas in which investigators should focus their efforts to provide a deeper understanding of IBD research areas in which rapid advances and novel concepts can be made. In addition, the Future Directions and Methods for IBD original research articles and review articles should include a discussion of areas in which improved methodologic tools have been identified and exactly how those methods should be performed. Basic science directions and methods should be presented that will expand our knowledge of areas that will allow novel insights to be made regarding the genetic, immunologic, microbial and environmental interactions that are the basis of the pathogenesis of Crohn's disease and ulcerative colitis. Clinical directions and methods should be presented that will allow investigators to make advances using cohort studies, multicenter registries, risk stratifications, and treatment outcomes. The goal of the Future Directions and Methods for clinical IBD original research articles and review articles should be to better understand the challenging biological variables in IBD patients and to provide optimal evidence of novel therapeutics that will more effectively treat, cure, and prevent Crohn's disease and ulcerative colitis.

Original research articles, for the Future Directions and Methods for IBD research section, should be written as described in the IBD Journal Instructions for original research articles. Review articles, for the Future Directions and Methods for IBD research section, should be no longer than 35 double spaced pages (including references and figure legends). There should be no more than 6 tables and figures (combined). Supplemental figures and tables will be allowed online. If page limits need to be increased, the authors may request permission from the Editors to increase the length of their future directions and methods for IBD research review article. The number of references should be limited to 100. The future directions and methods for IBD research review article should be focused on a single specific topic.

Letters to the Editor Guidelines Letters in response to articles published in the journal are welcome and should be submitted via IBD's online submission system. All Letters should start with the phrase "To the Editors," and be written as a letter. Letters must be submitted the end of the following calendar month (e.g. by the end of July, for letters referring to articles in the June print issue) and be a maximum of 400 words, with no more than one figure or table and no more than 5 references. All LTEs will be published online-only.

Research letters and case-based discussions will no longer be accepted for submission effective November 1, 2012.

REVISIONS

When submitting a revision, please submit both a clean copy and marked copy of the manuscript. The marked copy should highlight all of the changes made by authors after the original review. Authors can use the track changes feature of the Microsoft Word program to create a marked copy. Authors also should submit all tables and figures in separate files for production purposes.

AFTER ACCEPTANCE

Open access LWW's hybrid open access option is offered to authors whose articles have been accepted for publication. With this choice, articles are made freely available online immediately upon publication. Authors may take advantage of the open access option at the point of acceptance to ensure that this choice has no influence on the peer review and acceptance process. These articles are subject to the journal's standard peer-review process and will be accepted or rejected based on their own merit.

Authors of accepted peer-reviewed articles have the choice to pay a fee to allow perpetual unrestricted online access to their published article to readers globally, immediately upon publication. The article processing charge for *Inflammatory Bowel Diseases* is \$3,000. The article processing charge for authors funded by the Research Councils UK (RCUK) is \$3,800. The publication fee is charged on acceptance of the article and should be paid within 30 days by credit card by the author, funding agency or institution. Payment must be received in full for the article to be published open access.

- **Authors retain copyright** Authors retain their copyright for all articles they opt to publish open access. Authors grant LWW a license to publish the article and identify itself as the original publisher.
- **Creative Commons license** Articles opting for open access will be freely available to read, download and share from the time of publication. Articles are published under the terms of the Creative Commons License Attribution-NonCommercial No Derivative 3.0 which allows readers to disseminate and reuse the article, as well as share and reuse of the scientific material. It does not permit commercial exploitation or the creation of derivative works without specific permission. To view a copy of this license visit: <http://creativecommons.org/licenses/by-nc-nd/3.0>.
- **Compliance with NIH, RCUK, Wellcome Trust and other research funding agency accessibility requirements** A number of research funding agencies now require or request authors to submit the post-print (the article after peer review and acceptance but not the final published article) to a repository that is accessible online

by all without charge. As a service to our authors, LWW identifies to the National Library of Medicine (NLM) articles that require deposit and transmits the post-print of an article based on research funded in whole or in part by the National Institutes of Health, Howard Hughes Medical Institute, or other funding agencies to PubMed Central. The revised Copyright Transfer Agreement provides the mechanism. LWW ensures that authors can fully comply with the public access requirements of major funding bodies worldwide. Additionally, all authors who choose the open access option will have their final published article deposited into PubMed Central. RCUK and Wellcome funded authors can choose to publish their paper as open access with the payment of an article process charge (gold route), or opt for their accepted manuscript to be deposited (green route) into PMC with an embargo. With both the gold and green open access options, the author will continue to sign the Copyright Transfer Agreement (CTA) as it provides the mechanism for LWW to ensure that the author is fully compliant with the requirements. After signature of the CTA, the author will then sign a License to Publish where they will then own the copyright. Those authors who wish to publish their article via the gold route will be able to publish under the terms of the Attribution 3.0 (CCBY) License. To view a copy of this license visit: <http://creativecommons.org/licenses/by/2.0/>. Those authors who wish to publish their article via the green route will be able to publish under the terms of the Attribution Non-commercial 3.0 (CCBY NC) license (<http://creativecommons.org/licenses/by-nc/2.0/>). It is the responsibility of the author to inform the Editorial Office and/or LWW that they have RCUK funding. LWW will not be held responsible for retroactive deposits to PMC if the author has not completed the proper forms.

FAQ for open access <http://links.lww.com/LWW-ES/A48>

Page Proofs and Corrections - Corresponding authors will receive an e-mail containing a link to the electronic page proofs to check the copyedited and typeset article before publication. The page proofs are provided as portable document format (PDF) files which require Adobe Reader to be viewed and edited. Complete instructions will be provided with the e-mail for downloading the files and for returning the corrected pages electronically to the publisher. It is the author's responsibility to ensure that there are no errors in the proofs. Changes that have been made to conform to journal style should be allowed to stand if they do not alter the authors' meaning. Authors may be charged for alterations to the proofs beyond those required to correct errors or to answer queries. Proofs must be checked carefully and corrections returned within 24 to 48 hours of receipt, as requested in the communication accompanying the page proofs.

Reprints - The corresponding author will receive the reprint order form via e-mail while the article is in production. Reprint requests can be submitted any time before or after publication of the article. For orders submitted before the article is published, the reprints are normally shipped 6 to 8 weeks after publication of the issue in which the item appears. For any questions regarding reprints or publication fees, contact the Reprint Department by mail at Lippincott Williams & Wilkins, 351 West Camden Street, Baltimore, MD 21201; by phone at 1-800-341-2258; by fax at 410-528-4434; or by e-mail at: reprints@wolterskluwer.com.

Permissions - For permission and/or rights to use content for which the copyright holder is LWW or the society, please go to the journal's website and after clicking on the relevant article, click on the "Request Permissions" link under the "Article Tools" box that appears on the right side of the page. Alternatively, send an e-mail to customercare@copyright.com.

For Translation Rights & Licensing queries, contact Silvia Serra, Translations Rights, Licensing & Permissions Manager, Wolters Kluwer Health (Medical Research) Ltd, 250 Waterloo Road, London SE1 8RD, UK. Phone: +44 (0) 207 981 0600. E-mail: silvia.serra@wolterskluwer.com

For Special Projects and Reprints (U.S./Canada), contact Alan Moore, Director of Sales, Lippincott Williams & Wilkins, Two Commerce Square, 2001 Market Street, Philadelphia, PA 19103. Phone: 215-521-8638. E-mail: alan.moore@wolterskluwer.com

For Special Projects and Reprints (non-U.S./Canada), contact Silvia Serra, Translations Rights, Licensing & Permissions Manager, Wolters Kluwer Health (Medical Research) Ltd, 250 Waterloo Road, London SE1 8RD, UK. Phone: +44 (0) 207 981 0600. E-mail: silvia.serra@wolterskluwer.com

Agradecimentos

Ao meu orientador, Professor Doutor Fernando Magro, pela oportunidade de trabalhar com ele e pela disponibilidade ao longo de todo o trabalho.

À Eng. Joana Afonso, pelo apoio incansável e por todo o carinho.